



Poa labillardieri

Common Tussock-grass



Adult White-banded Grass-dart

Adult male (left) and female (right) Two-spotted Grass-skipper

(Greig, D, Australian Plant Image Index) (Allison, L, Flora of Westgate Park) (Grund, R, SA Butterflies & Moths)

(Grund, R, SA Butterflies & Moths)

Form: *Sedge or Flax* **Height:** 0.5 – 1.3 m **Spread:** 0.3 – 0.7 m **Flowers:** Green

Type: Spikes

Flowering time: Spring, Summer

Ecological functions:



Soils: sand, clay or loam; acidic, alkaline or neutral

Germination rate: 2 to 4 weeks

Seed collecting: place the drying seed inflorescence into a container and shake. Seeds and flower parts will fall out. Keep dry, use all these contents as your seed mix.

Seed treatment: none

Sowing requirements: sprinkle seeds over seed tray or broadcast over larger area, rake over and check that soil is lightly tilled. Press seeds in and keep moist. This species can also be propagated by division. Sow from Autumn.

Position: full sun, part shade.

Supplementary watering: minimal

Uses: an ornamental tufting perennial. Food source and habitat for small birds and lizards. Food host to White-banded Grass-dart (*Taractrocera papyria papyria*) and Two-spotted Grass-skipper (*Pasma tasmanica*) caterpillars.

Note: rejuvenate by trimming to just above ground level during late Winter and early Spring.